

# TOBACCO

Figure 1: Guide to interpreting spine plots




Table 1: Spine plot of smoking indicators. \* Prevalence of smoking among persons aged 16 years and over. \*\* Directly age-standardised rate per 100,000 population all ages. \*\*\* Directly age-standardised rate per 100,000 population aged 35 years and over. Data: London Health Observatory Local Tobacco Control Profiles for England

Indicator	Hounslow	London	England Worst	England Best		England Best
Smoking attributable deaths 2006 - 2008 ***	217.7	208.6	360.9	206.8		128.3
Smoking attributable deaths from heart disease 2006 - 2008 ***	33.2	33.7	62.4	34.0		18.6
Smoking attributable deaths from stroke 2006 - 2008 ***	8.7	9.4	18.9	9.6		5.2
Deaths from lung cancer 2006 - 2008 **	44.2	39.3	70.7	38.6		22.7
Deaths from COPD 2006 - 2008 **	29.3	27.7	52.3	26.6		15.4
Smokers during pregnancy 2008/2009	9.1	7.4	37.5	14.6		3.8
Estimated adult smoking prevalence 2009 - 2010 *	22.0	21.1	35.2	24.0		12.1
GP recorded smoking prevalence 2009/2010 *	14.6	16.8	33.4	18.8		11.1
Successful quitters at 4 weeks 2009/2010 *	1430.0	827.7	1933.9	894.7		405.7
Successful quitters at 4 weeks (CO Validated) 2009/2010 *	629.0	568.1	51.4	614.2		1455.3
Completeness of NS- SEC recording by Stop Smoking Services 2009/2010	67.0	54.5	1.6	66.1		98.1
Prescribed Nicotine Replacement Therapy 2009/2010	1583.0	2189.8	143.5	2997.0		10886.6
Cost of smoking attributable hospital admissions 2008/2009 ***	37.8	36.0	24.6	33.3		56.3
Smoking attributable hospital admissions 2008/2009 ***	1276.2	1258.3	2260.6	1265.3		794.4

Figure 2: Rate of successful quitters at 4-weeks per 100,000 smoking population aged 16 years and over: overall and CO validated




Figure 3: Percentage of adults who set a quit date with Stop Smoking Services by sex and age

